

Planning and Target-setting Form (Monthly)

ACTIVITIES AND BREAKDOWN OF MONTHLY TARGETS

RHU: _____

BHS: _____

| Activities | Timetable and Monthly Targets | | | | | | | | | | | | Annual Target | Resources Needed | Sources | |
|------------|---|---|---|---|---|---|---|---|---|---|---|---|---------------|------------------|---------|--|
| | J | F | M | A | M | J | J | A | S | O | N | D | | | | |
| A | TB AWARENESS CAMPAIGN | | | | | | | | | | | | | - | | |
| | Target Number of Households (HH) to be reached via: (list type of activities below) | | | | | | | | | | | | | | - | |
| | e.g., One-on-One | | | | | | | | | | | | | | | |
| | e.g., TB Caravan | | | | | | | | | | | | | | | |
| | e.g., Small-Group Discussion | | | | | | | | | | | | | | | |
| | e.g., Barangay Community Assembly | | | | | | | | | | | | | | | |
| | Total Target* | | | | | | | | | | | | | | | |
| B | CONDUCT OF CASE FINDING | | | | | | | | | | | | | | | |
| B.1 | Target for Presumptive TB Cases* | | | | | | | | | | | | | | | |
| | Target for TB cases* | | | | | | | | | | | | | | | |
| B.2 | Household Contacts | | | | | | | | | | | | | | - | |
| | Target 100% of household (HH) contacts of enrolled TB patients are screened | | | | | | | | | | | | | | - | |
| | Target 100% of HH contacts with symptoms are tested (DSSM/Xpert/PPD) | | | | | | | | | | | | | | | |
| | Target 100% of DSSM or Xpert positive enrolled to treatment | | | | | | | | | | | | | | | |
| | Target 100% of symptomatic DSSM-negative HH contacts undergo chest X-ray | | | | | | | | | | | | | | | |
| | Target 100% of symptomatic DSSM-negative HH contacts w/ CXR suggestive of TB enrolled to treatment | | | | | | | | | | | | | | | |
| B.3 | Cough Surveillance | | | | | | | | | | | | | | | |
| | 100% of households in CHV's catchment areas are screened for TB symptoms | | | | | | | | | | | | | | | |
| | 100% of household members with 2-week cough are tested (DSSM), and if DSSM-negative undergo chest X-ray | | | | | | | | | | | | | | | |
| | 100% of household members with other TB symptoms (not cough) are screened and tested | | | | | | | | | | | | | | | |
| B.4 | Household belonging to vulnerable population (DM, HIV) | | | | | | | | | | | | | | | |
| | 100% are referred for chest X-ray with or without symptoms | | | | | | | | | | | | | | | |
| B.5 | 100% of diagnosed TB cases are initiated treatment with treatment partners | | | | | | | | | | | | | | | |
| C | CASE HOLDING | | | | | | | | | | | | | | | |
| | 100% of enrolled TB cases have completed treatment | | | | | | | | | | | | | | | |

*From Planning and Target-setting Form (Annual)